

REPORT DOCUMENTATION PAGE			Form Approved OMB NO. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 5 January 99	3. REPORT TYPE AND DATES COVERED Final 10 Mar 95 - 30 Sep 98		
4. TITLE AND SUBTITLE Multivariate Approximation		5. FUNDING NUMBERS DAAH04-95-1-0089		
6. AUTHOR(S) Carl R. de Boor and Amos Ron				
7. PERFORMING ORGANIZATION NAMES(S) AND ADDRESS(ES) Center for the Mathematical Sciences University of Wisconsin-Madison 1308 W. Dayton St. Madison, WI 53715-1149		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211		10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARO 32830.19-MA		
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.		12 b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) Methods for representing multi-dimensional objects, such as functions of several variables and, more generally, (hyper-)surfaces is the main objective. One goal of such representation, whether approximate or exact, is the efficient evaluation of the object: <i>Multivariate Polynomial Interpolation</i> as well as <i>Scattered Data Approximation</i> both fall into this category. Another goal is a representation that allows one to identify and access easily and simultaneously relevant aspects of the object. The topics of <i>wavelets</i> and <i>Weyl-Heisenberg systems</i> belong to this second category. As to multivariate polynomial interpolation, we hope ultimately to duplicate the success of univariate polynomial interpolation as a basic tool in scientific computing. Current focus is the derivation of error formulae, using a multivariate divided difference just developed. Scattered data approximation is to be accomplished by extending well-known and efficient techniques for fitting to data on uniform meshes. Concerning Wavelets and Weyl-Heisenberg systems, the goal is to make new inroads in these important areas by applying tools and techniques from Approximation Theory, particularly those developed during studies of Shift-Invariant Spaces, a centerpiece of our previous research.				
14. SUBJECT TERMS Approximation Theory, Wavelets		15. NUMBER IF PAGES 4		16. PRICE CODE
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

Multivariate Approximation

Final Report

Carl de Boor & Amos Ron
December 29, 1998

for the
U. S. Army Research Office
DAAH04-95-1-0089

Center for the Mathematical Sciences
University of Wisconsin-Madison

approved for public release;
distribution unlimited

19990616 169

The view, opinion and/or findings contained in this report are those of the authors and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

DTIC QUALITY INSPECTED 1

A. STATEMENT OF THE PROBLEM STUDIED

Methods for representing multi-dimensional objects, such as functions of several variables and, more generally, (hyper-)surfaces is the main objective. One goal of such representation, whether approximate or exact, is the efficient evaluation of the object: *Multivariate Polynomial Interpolation* as well as *Scattered Data Approximation* both fall into this category. Another goal is a representation that allows one to identify and access easily and simultaneously relevant aspects of the object. The topics of *wavelets* and *Weyl-Heisenberg systems* belong to this second category.

As to multivariate polynomial interpolation, we hope ultimately to duplicate the success of univariate polynomial interpolation as a basic tool in scientific computing. Current focus is the derivation of error formulæ, using a multivariate divided difference just developed.

Scattered data approximation is to be accomplished by extending well-known and efficient techniques for fitting to data on uniform meshes.

Concerning Wavelets and Weyl-Heisenberg systems, the goal is to make new inroads in these important areas by applying tools and techniques from Approximation Theory, particularly those developed during studies of Shift-Invariant Spaces, a centerpiece of our previous research.

B. SUMMARY OF THE MOST IMPORTANT RESULTS

This brief outline relies on the fact that more details can be found in the (semi-)annual reports submitted during the grant period, and, if need be, in the manuscripts filed with ARO as required. All bracketed numbers refer to items in the list of publications given in C and in the Bibliography.

A new theory for *wavelet frames* was developed in [13]. The theory is based on tools and observations that were developed in the context of *shift-invariant spaces*, and in particular the 'fiberization' of the basic operators associated with a shift-invariant system, [20], [22] (as part of work on an earlier grant). The new theory of wavelet frames is based on the interplay between the given wavelet system, and a new, closely related, shift-invariant system: a *quasi-wavelet system*. The theory led to new algorithms for constructing wavelet frames from Multiresolution Analysis. The highlight of these algorithms is that there is essentially no restriction in them on the scaling function. As a result, compactly supported spline tight frames were constructed (in one and in several variables) [16]. The new frames may be useful for various wavelet applications, most notably in the areas of feature detection and noise removal. A study of the application of the aforementioned theory was initiated during the duration of this grant.

Several other studies in the interface between Approximation Theory and Wavelets were successfully completed. Of note is the theory of *approximation orders of finitely generated shift-invariant (FSI) spaces* (cf. [5]). The theory unravels some of the previous 'mysteries' in the studies of wavelets (such as the so-called 'sum-rules'), and was applied in the search, [10], of new factorization techniques of wavelet masks in one dimension. The connection between the approximation orders of wavelets and their smoothness was also studied, [11].

A better understanding of the error formula in the proposers' least polynomial interpolant was reached for special cases (see [4]), and a matlab code for the construction and evaluation of the least interpolant was extended to arbitrary dimensions and made available on the web (see http://www.cs.wisc.edu/~deboor/multiint/m_files.html).

Substantial progress has been made in the area of *approximation to scattered data*. First, a new method was developed, [21], that allows the conversion of the (readily available, and very well-understood) approximation schemes on uniform grids to schemes suitable for scattered data approximation. A concrete algorithm based on the general conversion method was developed in the Ph.D. thesis of **Jungho Yoon**. The algorithm deals with approximation to scattered data on bounded planar domains. Numerical experiments with a Matlab code developed for that purpose showed that the algorithm performs at least as well as, and in many instances better than, the state-of-the-art methods for scattered data approximation.

Some progress was made on the problem of proper parametrization of curves for use in the construction of surfaces by blending, in that **Scott Kersey** completed a thesis on the construction of a smoothest curve that fits, within given tolerances, to a given point sequence in any number of dimensions, with the smoothness measure being minimized also with respect to the parametrization used.

C. LIST OF ALL PUBLICATIONS AND TECHNICAL REPORTS

- [1] C. DE BOOR, *On the Sauer-Xu formula for the error in multivariate polynomial interpolation*, Math. Comp., 65 (1996), pp. 1231–1234.
- [2] C. DE BOOR, *On the Meir/Sharma/Hall/Meyer analysis of the spline interpolation error*, Approximation Theory and Optimization (M. D. Buhmann and A. Iserles, eds), Cambridge University Press, Cambridge, England, 1997, pp. 47–58.
- [3] C. DE BOOR, *The multiplicity of a spline zero*, Annals of Numerical Mathematics, 4 (1997), pp. 229–238.
- [4] C. DE BOOR, *The error in polynomial tensor-product, and in Chung-Yao, interpolation*, Surface Fitting and Multiresolution Methods (A. Le Méhauté, C. Rabut, and L. L. Schumaker, eds), Vanderbilt University Press, Nashville TN, 1997, pp. 35–50.
- [5] C. DE BOOR, R. DEVORE, AND A. RON, *Approximation orders of FSI spaces in $L_2(\mathbb{R}^d)$* , Constr. Approx., 14 (1998), pp. 631–652.
- [6] C. DE BOOR, N. DYN, AND A. RON, *On interpolation to data on linear manifolds in \mathbb{R}^d* , ms, 1993.
- [7] MARTIN D. BUHMANN, CHARLES A. MICCHELLI, AND AMOS RON, *Asymptotically optimal approximation and numerical solutions of differential equations*, Approximation Theory and Optimization (M. D. Buhmann and A. Iserles, eds), Cambridge University Press, Cambridge, England, 1997, pp. 59–82.
- [8] A. COHEN, I. DAUBECHIES, AND A. RON, *How smooth is the smoothest function in a refinable space (a note)*, Appl. Comput. Harmonic Anal., 3 (1996), pp. 87–89.
- [9] KARLHEINZ GRÖCHENIG AND AMOS RON, *Tight compactly supported wavelet frames of arbitrarily high smoothness*, Proc. Amer. Math. Soc., 126(4) (1998), pp. 1101–1107.

- [10] G. PLONKA AND A. RON, *A new factorization technique of the matrix mask of univariate refinable functions*, submitted, 1997.
- [11] A. RON, *Smooth refinable functions provide good approximation orders*, SIAM J. Math. Anal., 28 (1997), pp. 731–748.
- [12] A. RON AND ZUOWEI SHEN, *Gramian analysis of affine bases and affine frames*, Approximation Theory VIII, Vol. 2: Wavelets (Charles K. Chui and Larry L. Schumaker, eds), World Scientific Publishing Co., Inc., Singapore, 1995, pp. xxx–xxx.
- [13] A. RON AND ZUOWEI SHEN, *Affine systems in $L_2(\mathbb{R}^d)$: the analysis of the analysis operator*, J. Funct. Anal., 148 (1997), pp. 408–447.
- [14] AMOS RON AND ZUOWEI SHEN, *Affine systems in $L_2(\mathbb{R}^d)$ II: dual systems*, J. Fourier Anal. Appl., 3 (1997), pp. 617–637.
- [15] A. RON AND ZUOWEI SHEN, *Weyl-Heisenberg frames and Riesz bases in $L_2(\mathbb{R}^d)$* , Duke Math. J., 89 (1997), pp. 237–282.
- [16] A. RON AND ZUOWEI SHEN, *Compactly supported tight affine spline frames in $L_2(\mathbb{R}^d)$* , Math. Comp., 67 (1998), pp. 191–207.
- [17] A. RON AND ZUOWEI SHEN, *Construction of compactly supported affine frames in $L_2(\mathbb{R}^d)$* , Advances in Wavelets (K. S. Lau, ed), Springer-Verlag, New York, 1998, pp. xxx–xxx.
- [18] A. RON AND ZUOWEI ZHEN, *The Sobolev regularity of refinable functions*, submitted, 1997.
- [19] A. RON AND XINGPING SUN, *Strictly positive definite functions on spheres*, Math. Comp., 65(216) (1996), pp. 1513–1530.

**D. LIST OF ALL PARTICIPATING SCIENTIFIC PERSONNEL SHOWING
ANY ADVANCED DEGREES EARNED BY THEM
WHILE EMPLOYED ON THE PROJECT**

C. de Boor (entire time), A. Ron (entire time).

Two students were supported occasionally. Scott Kersey will officially finish in Spring'99, but has already successfully defended his thesis on "A minimizing spline curve under near-interpolatory constraints". Jungho Yoon finished in Spring'98, with a thesis on "Approximation in $L_p(\mathbb{R}^d)$ from a space spanned by the scattered shifts of radial basis function".

BIBLIOGRAPHY

- [20] C. DE BOOR, R. DEVORE, AND A. RON, *The structure of finitely generated shift-invariant spaces in $L_2(\mathbb{R}^d)$* , J. Funct. Anal., 119(1) (1994), pp. 37–78.
- [21] N. DYN AND A. RON, *Radial basis function approximation: from gridded centers to scattered centers*, Proc. London Math. Soc., 71(3) (1995), pp. 76–108.
- [22] A. RON AND ZUOWEI SHEN, *Frames and stable bases for shift-invariant subspaces of $L_2(\mathbb{R}^d)$* , Canad. J. Math., 47(5) (1995), pp. 1051–1094.